


Lebanon Pro



BRAND

FERTILIZER WITH Prodiamine 0.43% PREEMERGENCE WEED CONTROL

16-0-5



WITH
LSN[®]

LebanonTurf Stabilized Nitrogen

GUARANTEED ANALYSIS

Total Nitrogen (N)..... 16.0%
16.0% Urea Nitrogen*
Soluble Potash (K₂O)..... 5.0%

Derived from: Urea, Muriate of Potash.
Chlorine (Cl) not more than..... 5.0%
*8.0% Stabilized Urea Nitrogen N-(n-butyl) thiophosphoric acid.



F699

22-40158

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

SUGGESTED SPREADER SETTINGS			
Spreader	5.3#/1,000 sq. ft.	Spreader	5.3#/1,000 sq. ft.
LebanonTurf	5¼	Lesco (letter/Numeric)	K/22
Andersons AccuPro	0	PennMulch® HVO	N
Earthway Rotary	18	ProScape® SS	N
Gandy	30	Spyker	5¼
Lely	6½/l	Vicon (all models)	30

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

For selective preemergence control of grass and broadleaf weeds as listed in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Prodiamine..... 0.43%

OTHER INGREDIENTS: 99.57%

TOTAL 100.00%

EPA Reg. No. 961-362 (L) (D)
EPA Est. No. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

ProScape®, PennMulch® and LSN® LebanonTurf Stabilized Nitrogen are registered trademarks of Lebanon Seaboard Corporation.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Caution

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 888-208-1368 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.
www.LebanonTurf.com

Net Weight 50 LBS. (22.7 kg)

FERTILIZER

WITH

Prodiamine 0.43%

PREEMERGENCE WEED CONTROL

16-0-5




WITH **LSN**®

LebanonTurf Stabilized Nitrogen

GUARANTEED ANALYSIS

Total Nitrogen (N).....	16.0%
16.0% Urea Nitrogen*	
Soluble Potash (K ₂ O).....	5.0%
Derived from: Urea, Muriate of Potash.	
Chlorine (Cl) not more than.....	5.0%
*8.0% Stabilized Urea Nitrogen N-(n-butyl) thiophosphoric acid.	



F699

22-40158

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

SUGGESTED SPREADER SETTINGS

Spreader	5.3#/1,000 sq. ft.	Spreader	5.3#/1,000 sq. ft.
LebanonTurf	5¼	Lesco (letter/Numeric)	K/22
Andersons AccuPro	0	PennMulch® HVO	N
Earthway Rotary	18	ProScape® SS	N
Gandy	30	Spyker	5¼
Lely	6½II	Vicon (all models)	30

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

For selective preemergence control of grass and broadleaf weeds as listed in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:

Prodiamine.....	0.43%
OTHER INGREDIENTS:	99.57%
TOTAL	<u>100.00%</u>

EPA Reg. No. 961-362

EPA Est. No. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

ProScape®, PennMulch® and LSN® LebanonTurf Stabilized Nitrogen are registered trademarks of Lebanon Seaboard Corporation.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Precautionary Statements
Hazards to Humans and Domestic Animals
Caution

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

- IF IN EYES:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- IF ON SKIN OR CLOTHING:
- Call a poison control center or doctor for treatment advice.
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- IF INHALED:
- Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
You may also contact 888-208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com

Net Weight 50 LBS. (22.7 kg)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many listed grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aeri ally.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS


ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses and lawns. The maximum amount of this product that may be applied per year is given for each turfgrass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED

GERMINATION DATES



Approximate Date



After May 30



After May 10



After April 20



After March 20



January 1 to March 20

USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection must be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is for use on the turfgrass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of LEBANON PRO BRAND FERTILZIER WITH PRODIAMINE 0.43% PREEMERGENCE WEED CONTROL by Turfgrass Species¹⁾

Turf Species:	Lbs. Product/Acre	Lbs. AI/Acre	Lbs. Product/1,000 Sq. Ft.
Bermudagrass ²⁾ , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	349	1.5	8.0
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	233	1.0	5.3
Creeping Red Fescue	175	0.75	4.0
Creeping Bentgrass	150	0.65	3.4

¹⁾ These are the maximum rates per calendar year by species limitations.

²⁾ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 121 lbs./Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescue grass due to reduced product rates used in sprigging situations.

- Do not apply more than 349 lbs. of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Chickweed, Mouseear (from seed)	Foxtails, Annual	Johnsongrass (from seed)	Lambsquarter, Common	Purslane, Common	Speedwell, Persian
Bluegrass, Annual (<i>Poa annua</i>)	Crabgrass (large, smooth)	Goosegrass ¹	Junglerice	Lovegrass	Pusley, Florida	Sprangletop
Carpetweed	Crowfootgrass	Henbit	Knotweed	Panicum (Texas, Fall, Browntop)	Rescuegrass ³	Spurge, Prostrate
Chickweed, Common	Cupgrass, Woolly	Itchgrass	Kochia	Pigweed	Shepherd's Purse ²	Witchgrass
					Signalgrass, Broadleaf	Woodsorrel, Yellow (from seed)

¹ In many areas a single application of 150 to 349 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 150 to 233 lbs./Acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rate Table**. Do not exceed the maximum rate for turfgrass species listed in the **Maximum Annual Rate Table** above.

² Apply this product in late summer, fall, or winter prior to germination.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

LEBANON PRO BRAND FERTILZIER WITH PRODIAMINE 0.43%
PREEMERGENCE WEED CONTROL Rate (lb./A)

Product	AI	Months Before Overseeding		
		North	Transition	South
116	.50	4	4	4
150	.66	5	4	4
175	.74	6	5	5
185	.80	---	6	6
233	1.00	---	7	7
265	1.14	---	---	9
302	1.30	---	---	10
349	1.50	---	---	12

LANDSCAPE ORNAMENTALS (including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 150 to 349 lbs./Acre in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications are to be made so long as the total amount of product applied does not exceed the maximum annual application rate of 349 lbs./Acre.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to established ornamentals as a broadcast, over the top or directed application.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

Apply this product in the fall and/or spring to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. Apply this product over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Fir species** (Balsam, Fraser, Noble, etc.)	Wild Lilac <i>Ceratostigma plumbagonoides</i>	Gladiolus species**	Glossy Privet (Waxleaf)	Monterey Pine	Pincushion Flower
Abelia: Sherwood	False Cypress	Baby's Breath	Lily: Jazz	Japanese Black Pine	Stonecrop (<i>thalictum dipterocarpum</i>)
Japanese Maple	<i>Chrysanthemum nipponicum</i>	English Ivy	Big Blue Lily; Turf; Silver Mound;	Eastern White Pine	Stonecrop; Lidakense
Norway Maple	Cleyera	Sunrose	Evergreen Giant	Scotch Pine	Stonecrop (<i>truchelspermum asiatum</i>)
Yarrow: King Edward	Citrus species*	Daylily: Aztec Gold, Stella De Oro,	Liriope, Variegated	Loblolly Pine	Stonecrop; Dragon's Blood
Kiwi*	Coreopsis (Calliopsis): Early	Tender Love	Liriope, Creeping	Virginia Pine	Spiraea: Anthony Waterer
Lily-of-the-Nile (African Lily)	Sunrise, Moonbeam	Rose of Sharon**	Cardinal Flower, Indian Pink	Pistachio*	Australian Brushcherry
<i>Agapanthus orientalis</i>	Flowering Dogwood	Mallow: Disco Belle White	Japanese Honeysuckle	Queensland Pittosporum	Japanese Yew
Five-Leaf or Chocolate Vine	American Dogwood	Chinese Hibiscus	Tatarian Honeysuckle	Japanese Pittosporum	Yew
Lady's Leek, Nodding Onion	Pampas Grass	Hosta, Plantain Lily (Fragrant)	Burgundy	Japanese Yew	Germander
Japanese Anemone	Cranberry Cotoneaster	Hosta, "Searsucker"	Loosestrife; Modern Pink	English Laurel	Meadow Rue
Aquilegia: Red and Gold	Cotoneaster	<i>Houttuynia cordata var. variegata</i>	Magnolia species**	Almond, Apricot, Nectarine, Peach, Plum	American Arborvitae
Vine Hill Manzanita	Bearberry Cotoneaster	Bigleaf Hydrangea	Ice Plant	and Prune*	Star Jasmine
Cape Weed	Rockspray Cotoneaster	Holly	Crabapple*	Douglas Fir**	Canada Hemlock
Wormwood; Silver Mound, Castle	Hawthorne	Chinese Holly	Yaku Jima, Silberfeder	Firethorn (Scarlet)	Tulip species
Aster: Bonny Blue, Purple Dome	Lucifer	Japanese Holly; Helleri	Heavenly Bamboo	Firethorn (<i>P. Fortuneana</i>)	Veronica, Speedwell; Sunny Border
<i>Aster X Frikartii</i>	Italian Cypress	American Holly	Narcissus species**	Firethorn (<i>P. Koidzumii</i>)	Japanese Viburnum
Lady Fern; Fern Lady	White Trailing Ice Plant	Yaupon Holly; Schillings	Oleander	Bradford Pear sp.	Sweet Viburnum
Japanese Aucuba	Cooperi Pink	<i>Inula ensifolia</i>	Olive*	Oak, Shumard's Red	Japanese Snowball
Fibrous Begonia: Hardy Grandis	Larkspur: Blue Elf	Sword-Leaved Iris; Jodlesong	Mondo Grass	Oak species	Canary Island Viburnum
Barberry	Cheeddar Pink	Siberian Iris; Cabernet	Osmanthus (False Holly): Gulf Tide	Indian Hawthorne	Arrowwood Viburnum
Wintergreen Barberry	Hop Bush	Iris species**	Trailing African Daisy	Yedda Hawthorn	Laurustinus
Mentor Barberry	Coneflower, Purple; Magnus	Winter Jasmine	Sourwood	'Coral Bells'	Cranberry Bush
Japanese Barberry	Silverberry	Chinese Juniper; Nick's Compact,	Tree Peony	'Delaware Valley White'	Leatherleaf Viburnum
Warty Barberry	Wintercreeper	Parsonii	Fountain Grass (Dwarf)	'Flame Creeper'	Greater Periwinkle, Vinca
<i>Bergenia cordifolia</i>	Japanese Spindle Tree (Evergreen Euonymus)	Shore Juniper; Blue Pacific	<i>Perovskia atriplicifolia</i>	'Formosa'	Common Periwinkle, Vinca
Snowbank	Spreading Euonymus	Parsoni	Avocado*	'Girard Crimson'	Grape*
Butterfly-Bush (Dwarf Blue);	Japanese Aralia	Creeping Juniper	Frasier's Photinia (Redtip)	'George L. Tabor'	Old Fashioned Weigela
Royal Red	Border Forsythia	Walnut*	Dragonhead, False; Vivid	'Hino-crimson'	Spanish Bayonet, Yucca
Japanese Boxwood	Weeping Forsythia	Shrimp Plant	Spruce species** (Colorado Blue,	'PJM'	Adam's Needle, Yucca
Weeping Bottlebrush	Greenstem Forsythia	Crape Myrtle	Norway, etc.)	'Roseum Elegans'	
Scotch Heather	Gaillardia, Blanket Flower: "Goblin"	Crape Myrtle; Tuscarora	Lily-of-the-Valley Bush	'Wakeiebisu'	
Tussock Bellflower, (White Clips)	Gardenia, Cape-Jasmine, Radicans	Weeping Lantana	Calabrian Pine	'White Gumbo'	
Trumpet Creeper, Trumpet Flower,	<i>Gaura sp.</i>	Lavender; Munstead	Canary Island Pine	Lady Bank's Rose	
Madame Galen	Gentian	Edelweiss	Slash Pine	Rosemary*	
Hottentot Fig (Ice Plant)	Cranesbill	Chinese Privet; Variegata	Aleppo Pine	Black-Eyed Susan; Goldstrum	
Feathery Cassia		Amur Privet	Austrian Pine	Santolina	
		Japanese Privet	Longleaf Pine	Saxifrage; Purple Dome	

* Do not use on food producing trees, vines, or plants. ** Not for use on container grown plants.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.